



February 2015

*This fact sheet is a quick reference on demographics and the use of services by people with spinal cord injury (SCI).*

*The National SCI Database is a prospective longitudinal multicenter study that captures data from an estimated 13% of new SCI cases in the U.S. from 28 federally funded SCI Model Systems since 1973. The database has demographic and condition status data through 2014 for 30,532 people with SCI.*

Incidence and prevalence statistics are estimates obtained from several studies. These statistics are not derived from the National SCI Database. All the remaining statistics in this fact sheet are solely based on data from the National SCI Database. Page 1 statistics focus on the new injuries that occurred since 2010. Page 2 statistics are about long-term outcomes, analyzing baseline and follow-up data from all database participants since 1973.

Presently, there are 14 systems and 5 Form II centers sponsored by the National Institute on Disability and Rehabilitation Research, Office of Special Education and Rehabilitative Services, U.S. Department of Education. For a complete list, please visit <http://www.msktc.org/sci/model-system-centers>.

**The National SCI Statistical Center**  
1717 6th Avenue South, SRC 515,  
Birmingham, AL 35233-7330

*For Statistics: (205) 934-3342;  
For Business: (205) 934-3320;  
TDD: (205) 934-4642;  
FAX: (205) 934-2709;  
E-mail: [NSCISC@uab.edu](mailto:NSCISC@uab.edu)*

Website:  
<https://www.nscisc.uab.edu/>

## Incidence

It is estimated that the annual incidence of SCI, not including those who die at the scene of the accident, is approximately 40 cases per million population in the U.S. or approximately 12,500 new cases each year, given the current population size of 313 million people in the U.S.

## Prevalence

The number of people in the U.S. who are alive in 2014 who have SCI has been estimated to be approximately 276,000 persons, with a range from 240,000 to 337,000 persons.

## Age at Injury


The average age at injury has increased from 29 years during 1970s to 42 years since 2010.

## Gender

Approximately 80% of spinal cord injuries occur among males.

## Race/Ethnicity

Currently, about 23% of spinal cord injuries occur among blacks, which is higher than the proportion of blacks in the general population (12%).



## Etiology


Vehicle crashes are the leading cause of injury, followed by falls, acts of violence (primarily gunshot wounds), and sports.

## Length of stay

Days hospitalized in the acute care unit have declined from 24 days in the 1970s to 11 days since 2010. Substantial downward trends are also noted for days in the rehabilitation unit (from 98 to 36 days).

## Neurologic level and extent of lesion

The most frequent neurologic category is incomplete tetraplegia followed by incomplete paraplegia, complete paraplegia, and complete tetraplegia. Less than 1% of persons experienced complete neurologic recovery by hospital discharge.



## Marital status

More than half of persons with SCI are single/never married when injured. The percentage of persons who are married slowly increases as the years post-injury increase, as does the divorce rate.

Status (%)	At injury	Year 1	Year 10	Year 20	Year 30	Year 40
Single	51.6	50.4	41.6	35.1	29.3	14.3
Married	32.7	32.0	33.6	36.0	39.4	42.9
Divorced	9.5	11.2	19.2	23.8	24.7	17.9

## Occupational status

At one year after injury, 12% of persons with SCI are employed, and by 20 years post-injury, about one third are employed.

Status (%)	At injury	Year 1	Year 10	Year 20	Year 30	Year 40
Employed	58.1	12.2	27.9	34.4	32.9	18.5
Student	15.3	16.1	7.2	2.7	0.7	0.0

## Lifetime costs

The average yearly health care and living expenses and the estimated lifetime costs that are directly attributable to SCI vary greatly according to severity of injury. These figures do not include any indirect costs—such as losses in wages, fringe benefits, and productivity, which average \$71,961 per year in 2014 dollars—but vary substantially based on education, severity of injury, and pre-injury employment history.

Severity of Injury	Average Yearly Expenses (in 2014 dollars)		Estimated Lifetime Costs by Age At Injury (discounted at 2%)	
	First Year	Each Subsequent Year	25 years old	50 years old
High Tetraplegia (C1–C4) AIS ABC	\$1,064,716	\$184,891	\$4,724,181	\$2,596,329
Low Tetraplegia (C5–C8) AIS ABC	\$769,351	\$113,423	\$3,451,781	\$2,123,154
Paraplegia AIS ABC	\$518,904	\$68,739	\$2,310,104	\$1,516,052
Motor Functional at Any Level AIS D	\$347,484	\$42,206	\$1,578,274	\$1,113,990

Data Source: Economic Impact of SCI published in the journal *Topics in Spinal Cord Injury Rehabilitation*, Volume 16, Number 4, in 2011.

**Life expectancy** is the average remaining years of life for an individual. Life expectancies for persons with SCI are still significantly below life expectancies for those without SCI and have not improved since the 1980s. Mortality rates are significantly higher during the first year after injury than during subsequent years, particularly for severely injured persons.

Age at Injury	Life expectancy (years) for post-injury by severity of injury and age at injury										
	For persons who survive the first 24 hours					For persons surviving at least 1 year post-injury					
	No SCI	AIS D—Motor Functional at Any Level	Para	Low Tetra (C5–C8)	High Tetra (C1–C4)	Ventilator Dependent Any Level	AIS D—Motor Functional at Any Level	Para	Low Tetra (C5–C8)	High Tetra (C1–C4)	Ventilator Dependent Any Level
20	59.3	52.6	45.0	39.9	35.6	19.2	53.0	45.5	40.6	36.9	25.3
40	40.4	34.2	27.6	23.3	19.9	8.7	34.6	28.0	23.9	20.9	12.4
60	23.0	18.0	13.0	10.1	8.0	2.1	18.2	13.3	10.5	8.6	3.9

**Cause of death:** Persons enrolled in the National SCI Database since its inception in 1973 have now been followed for 40 years after injury. During that time, the causes of death that appear to have the greatest impact on reduced life expectancy for this population are pneumonia and septicemia. Mortality rates are declining for cancer, heart disease, stroke, arterial diseases, pulmonary embolus, urinary diseases, digestive diseases, and suicide. However, these gains are being offset by increasing mortality rates for endocrine, metabolic and nutritional diseases, accidents, nervous system diseases, musculoskeletal disorders and mental disorders. There has been no change in the mortality rate for septicemia in the past 40 years, and only slight improvement in mortality due to respiratory diseases.

This is a publication of the National Spinal Cord Injury Statistical Center, Birmingham, Alabama, (Grant Number H133A110002), and the Model Systems Knowledge Translation Center at American Institutes for Research, Washington, DC (Grant Number H133A110004). Both are funded by the National Institute on Disability and Rehabilitation Research, Office of Special Education and Rehabilitative Services, U.S. Department of Education, Washington, DC. The opinions contained in this publication are those of the grantee and do not necessarily reflect those of the U.S. Department of Education. © 2015, Board of Trustees, University of Alabama.

Citation: National Spinal Cord Injury Statistical Center, Facts and Figures at a Glance. Birmingham, AL: University of Alabama at Birmingham, February 2015.

## Education

Over half of persons with SCI are high school graduates at time of injury. Level of education slowly increases over post-injury years.

Education (%)	At injury	Year 1	Year 10	Year 20	Year 30	Year 40
High school only	51.5	54.1	51.0	46.4	44.4	33.3
College or higher	10.6	11.8	21.7	29.3	35.9	40.7

## Re-hospitalization

About 30% of persons with SCI experience one or more hospitalizations during a 12-month period. Among those re-hospitalized the length of hospital stay averages about 22 days. Diseases of the genitourinary system are the leading cause of re-hospitalization, followed by disease of the skin. Respiratory, digestive, circulatory, and musculoskeletal diseases are also common causes.